THIRD PARTY MANUFACTURING ALSO AVAILABLE FOR ALL OTHER PHARMACEUTICAL PRODUCTS



F9 DRUGS & CO. GMP CERTIFIED COMPANY

A trusted manufacturer & exporter of high-quality veterinary medicines, including oral liquids and tablets with Quality and safety

Contact us For Custom Packaging or Third Party Manufacturing DEHRADUN, INDIA-248001 Email : furever9@gmail.com || M. +91 8920240034



Exporting Medicine Worldwide

FOLLOW US ON





www.f9drugs.com



furever9@gmail.com

VETERINARY ORAL MEDICINE LIQUID	
ENROFUR	ENROFLOXACIN 20% ORAL SUSPENSION
IVOFUR	IVERMECTIN ORAL SUSPENSION



Exporting Medicine Worldwide

FOLLOW US ON 🗗 🙆 🎇

F9 DRUGS & CO.

ENROFUR

ENROFLOXACIN 20% ORAL SUSPENSION

COMPOSITION

Each ml contains Enrofloxacin BP Vet 200 mg

INDICATIONS

Enrofloxacin 20% Oral Solution is a broad-spectrum antimicrobial effective against a wide range of bacteria and mycoplasma.

Recommended for Poultry in the treatment of:

- Chronic Respiratory Disease (CRD)
- Fowl Cholera (Avian Hemorrhagic Septicemia)
- Infectious Coryza
- Salmonellosis
- lostridiosis (Necrotic Enteritis)
- Collibacillosis
- Omphalitis
- Other infections caused by *Enrofloxacin*-sensitive organisms

DOSAGE AND ADMINISTRATION

- General dose: 1 ml per 4 litres of drinking water for 3 days
 continuously
- For Salmonella and Mycoplasma infections: Continue for 5 days (Equivalent to 10 mg Enrofloxacin per kg body weight)

WITHDRAWAL PERIOD

Meat: 12 days



FUREVER 9 CORPORATION 207, Turner Road, Clement Town Dehradun 248001 (U.K.) India.

PACK SIZE Available in 100 ml & 200 ml ENROFU

ENROFLOXACIN 20%



COMPOSITION:

Each ml contains : Ivermectin 0.8 mg Excipients ad q.s.

DESCRIPTION:

IVOFUR LIQUID is an oral antiparasitic formulation containing Ivermectin, a broad-spectrum endectocide from the avermectin class. It is effective against a wide range of gastrointestinal roundworms and ectoparasites

INDICATIONS

For the treatment of:

- Gastrointestinal roundworms
- Lungworm
- Lice
- Oestriasis
- Scabies

Effective against:

• Trichostrongylus, Cooperia, Ostertagia, Haemonchus, Nematodirus, Chabertia, Bunostomum, Dictyocaulus spp. In Calves, Sheep, and Goats

CONTRAINDICATIONS

• Not to be administered to lactating • Meat: 14 days animals



FUREVER 9 CORPORATION 207, Turner Road, Clement Town Dehradun 248001 (U.K.) India.

VETERINARY / NOT FOR HUMAN USE / FOR ANIMAL TREATMENT ONLY

SIDE EFFECTS

- Musculoskeletal pain
- · Edema of the face or extremities
- · Itching and papular rash
- · Temporary, mild coughing posttreatment (not clinically significant)

DOSAGE AND

ADMINISTRATION: For oral use only

· General dose: 2.5 ml per 10 kg body weight

WITHDRAWAL PERIOD

PACK SIZE Available in 500 ml & 1 ltr



IVERMECTIN

LIQUID

VOFUR