

**THIRD PARTY MANUFACTURING ALSO AVAILABLE
FOR ALL OTHER PHARMACEUTICAL PRODUCTS**



DRUGS & CO.
www.f9drugs.com

F9 DRUGS & CO.

GMP CERTIFIED COMPANY

**A trusted manufacturer & exporter of high-quality veterinary medicines,
including oral liquids and tablets with Quality and safety**

Contact us For Custom Packaging or Third Party Manufacturing

DEHRADUN, INDIA-248001

Email : furever9@gmail.com || M. +91 8920240034



Exporting Medicine Worldwide

FOLLOW US ON



VETERINARY ORAL MEDICINE POWDER FOR POULTRY AND LIVE STOCK

AMOXIN 200	AMOXYCILLIN ORAL POWDER
AMOXIN C	AMOXYCILLIN AND COLISTIN ORAL POWDER
AMPROVIT S	AMPROLIUM AND SULFAQUINOXALINE POWDER WITH VIT A1 AND K3
BIOMIN VITA	OXYTETRACYCLINE AND NEOMYCIN WITH VITAMINS AND MILERALS
DEWORMIT F	FLUBENDAZOLE 5% ORAL POWDER
ERYCIN	ERYTHROMYCIN ORAL POWDER
EZYDOXIN 200	DOXYCYCLINE ORAL POWDER
EZYDOXIN C	DOXYCYCLINE AND COLISTIN ORAL POWDER
EZYDOXIN G	DOXYCYCLINE AND GENTAMICIN ORAL POWDER
GASTROFLU	FLUMEQUINE POWDER
HYDRODEX	DEXAMETHASONE AND HYDROCHLOROHIAZIDE ORAL POWDER
LEVASAFE	LEVAMISOLE HYDROCHLORIDE ORAL POWDER
METROFUR	METRONIDAZOLE 10% ORAL POWDER
NEOWIN	NEOMYCIN ORAL POWDER
OXYFUR 40%	OXYTETRACYCLINE 40% POWDER
TRIPower K3	SULPHADIMIDINE, SULPHADIAZINE, SULPHATHIAZOLE POWDER WITH VITAMIN K3
PEZINE	PIPERAZINE CITRATE ORAL POWDER
TYLODOX	TYLOSIN AND DOXYCYCLINE ORAL POWDER
TYLOSOL	TYLOSIN ORAL POWDER



Exporting Medicine Worldwide

FOLLOW US ON



F9 DRUGS & CO.

AMOXIN 200

AMOXYCILLIN TRIHYDRATE POWDER

COMPOSITION

Each gram contains :

Amoxycillin trihydrate 200 mg
Excipients ad 1 g

INDICATIONS

Indicated for the treatment of gastrointestinal, respiratory, and urinary tract infections caused by amoxycillin-sensitive microorganisms, including: *Campylobacter*, *Clostridium*, *Corynebacterium*, *Escherichia coli*, *Erysipelothrix*, *Haemophilus*, *Pasteurella*, *Salmonella*, penicillinase-negative *Staphylococcus spp.*, and *Streptococcus spp.* Applicable for use in **Calves, Goats, Poultry, Sheep, and Swine**.

CONTRAINDICATIONS

- Hypersensitivity to amoxycillin
- Use in animals with severely impaired renal function
- Concurrent use with tetracyclines, chloramphenicol, macrolides, or lincosamides
- Avoid in animals with active microbial digestion (e.g., mature ruminants)

DOSAGE & ADMINISTRATION

For oral administration:

- **Calves, Goats, Sheep:** 5 g per 100 kg body weight, twice daily for 3–5 days
- **Poultry, Swine:** 1 kg per 1000–2000 litres of drinking water for 3–5 days

Note: Intended for pre-ruminant calves, lambs, and kids only.


Or As directed by the Veterinarian.

WITHDRAWAL PERIOD

- **Meat:**
 - Calves, Goats, Sheep, Swine: 8 days
 - Poultry: 3 days



FUREVER 9 CORPORATION
207, Turner Road, Clement Town
Dehradun 248001 (U.K.) India.



VETERINARY
NOT FOR HUMAN USE
FOR ANIMAL TREATMENT ONLY



PACK SIZE
Available in
1 Kg Jar
100 g Pouch

AMOXIN C

Colistin Sulphate & Amoxycillin Trihydrate

COMPOSITION

Each gram contains :


Colistin sulphate 1 200 100 IU
Amoxycillin trihydrate 210 mg
Excipients ad 1 g

INDICATIONS

Indicated for the treatment of **gastrointestinal, respiratory, and urinary tract infections** caused by microorganisms sensitive to amoxycillin and colistin, including:

Campylobacter, Clostridium, Corynebacterium, Escherichia coli, Erysipelothrix, Haemophilus, Pasteurella, Salmonella, penicillinase-negative Staphylococcus spp., and Streptococcus spp.

Applicable for use in **Calves, Goats, Poultry, Sheep, and Swine.**



**VETERINARY
NOT FOR HUMAN USE
FOR ANIMAL TREATMENT ONLY**

CONTRAINDICATIONS

- Hypersensitivity to amoxycillin and/or colistin
- Use in animals with severely impaired renal function
- Concurrent administration with tetracyclines, chloramphenicol, macrolides, or lincosamides
- Avoid use in animals with active microbial digestion (e.g., mature ruminants)

DOSAGE AND ADMINISTRATION

For oral administration :

Calves, Sheep and Goats Preventive :

1 g per 50-100 kg body weight through drinking water or milk for 21 days.

Poultry : 20 g per 20-40 litres of drinking water for 5-7 days

Curative : 5 g per 25-50 kg body weight through drinking water or milk for 5 days.

Or As directed by the Veterinarian.

WITHDRAWAL PERIOD

- **Meat:** 8 days



FUREVER 9 CORPORATION
207, Turner Road, Clement Town
Dehradun 248001 (U.K.) India.



PACK SIZE
Available in
1 Kg Jar
100 g Pouch


AMPROVIT S

Powder For Oral Administration

COMPOSITION

Each 100 gram contains :

Amprolium Hydrochloride	17.5 g
Sulfaquinoxaline	17.5 g
Vitamin A1 retinol acetate	1,500,000 I.U
Vitamin K3	500 mg
Excipients	q.s.



INDICATIONS

Coccidiosis in Calves, Sheep, Goat and Poultry

DOSAGE AND ADMINISTRATION

For oral administration :

Calves, Sheep and Goats Preventive :

1 g per 50-100 kg body weight through drinking water or milk for 21 days.

Poultry : 20 g per 20-40 litres of drinking water for 5-7 days

Curative : 5 g per 25-50 kg body weight through drinking water or milk for 5 days.

WITHDRAWAL PERIOD

For meat :

Calves, Sheep and Goats : 14 days.

Poultry : 14 days.

VETERINARY
NOT FOR HUMAN USE
FOR ANIMAL TREATMENT ONLY



FUREVER 9 CORPORATION
207, Turner Road, Clement Town
Dehradun 248001 (U.K.) India.



PACK SIZE
Available in
1 Kg Jar
100 g Pouch

BIOMIN VITA

Antibiotics and Vitamins

COMPOSITION

Each gram contains :

Oxytetracycline hydrochloride	60 mg
Neomycin sulphate	40 mg
Vitamin A, retinol acetate	7500 IU
Vitamin D3, cholecalciferol	1500 IU
Vitamin E, α-tocopherolacetate	5 mg
Vitamin B1, thiamine hydrochloride	1 mg
Vitamin B2, riboflavin	2 mg
Vitamin B6, pyridoxine hydrochloride	2 mg
Vitamin B12, cyanocobalamin	7.5 mg
Vitamin C, ascorbic acid	25 mg
Ca-pantothenate	7.5 mg
Vitamin K3, menadione sodium bisulphite	5 mg
Nicotinamide	15 mg
Folic acid	0.3 mg
Methionine	30 mg
Lysine	50 mg
Excipients ad	1 g

INDICATIONS

BIOMIN VITA is a potent formulation combining broad-spectrum antibiotics with essential vitamins. It is designed to:

- Boost egg production
- Promote growth and enhance feed efficiency
- Serve as a vitamin supplement during disease outbreaks and stress


It is effective against gastrointestinal, respiratory, and urinary tract infections caused by organisms sensitive to oxytetracycline and neomycin. These include:

Bordetella, Campylobacter, Chlamydia, E. coli, Haemophilus, Klebsiella, Mycoplasma, Pasteurella, Rickettsia, Salmonella, Staphylococcus, and Streptococcus spp.

Applicable for use in Calves, Goats, Sheep, Poultry, and Swine.

CONTRAINDICATIONS

- Do not administer to animals hypersensitive to tetracyclines or aminoglycosides.
- Avoid use in animals with severely impaired liver or kidney function.
- Do not use concurrently with bactericidal antibiotics such as penicillins.
- Not recommended for animals with active microbial digestion (e.g., ruminants).



DOSAGE AND ADMINISTRATION

For oral use only:

- **Poultry and Swine**
 - **Prevention:** 1 kg per 2000 liters of drinking water for 5–7 days
 - **Treatment:** 1 kg per 1000 liters of drinking water for 5–7 days
- **Calves, Sheep, and Goats**
 - 1 g per 5 kg body weight daily for 5–7 days
 - *Note: Intended for pre-ruminant calves, lambs, and kids only*

**VETERINARY / NOT FOR HUMAN USE
FOR ANIMAL TREATMENT ONLY**

WITHDRAWAL PERIOD

- Meat: 7 days
- Eggs: 1 day



PACK SIZE
Available in
Sachets of 30 g and 100 g
Jars of 1000 g



FUREVER 9 CORPORATION
207, Turner Road, Clement Town
Dehradun 248001 (U.K.) India.

DEWORMIT-F

FLUBENDAZOLE POWDER 5%

COMPOSITION

Each 100 gram contains :

Flubendazole 5 g
Excipient q.s.

INDICATIONS

DEWORMIT-F 5% Powder is indicated for the prevention and treatment of the following parasitic infections:

- **Swine:** *Ascaris suum*, *Hyostrongylus rubidus*, *Oesophagostomum dentatum*, *Trichuris suis*, *Strongyloides ransomi*, *Metastrongylus apri*
- **Poultry (Chicken, Turkey, Pheasant):** *Syngamus trachea*, *Ascaridia galli*, *Heterakis gallinarum*, *Capillaria spp.*, *Amidostomum anseris*, *Trichostrongylus tenius*, *Railletina spp.*


SPECIAL PRECAUTIONS FOR USE

- For optimal results, maintain strict hygiene in pig pens and poultry housing during treatment.
- Periodic fecal examinations are recommended to verify treatment efficacy.
- In case of accidental ingestion, seek immediate medical attention and provide the physician with the product label or leaflet.
- Individuals sensitive to benzimidazoles should avoid handling this product.
- Do not eat, drink, or smoke while administering the product.

DOSAGE & ADMINISTRATION

DEWORMIT-F 5% Powder should be thoroughly mixed with feed prior to administration.

- **Pigs (Group Treatment):**
 - **Piglets and Fattening Pigs:** 30 ppm Flubendazole (equivalent to 600 g FUBACIN 5% per ton of feed) for 5 consecutive days.
 - For severe *Trichuris suis* infections, extend treatment to 10 days.
- **Breeding Pigs:** 30 ppm Flubendazole (600 g FUBACIN 5% per ton of feed) for 10 consecutive days.
- **Individual Treatment:** 5 mg/kg body weight of Flubendazole (10 g FUBACIN 5% per 100 kg body weight) mixed with or sprinkled over feed for 5 days.
- For severe *Trichuris suis* infections, extend treatment to 10 days.
- **Poultry:**
 - **Chickens:** 30 ppm Flubendazole (600 g DEWORMIT-F 5% per ton of feed) for 7 consecutive days.
 - For tapeworm (*Railletina spp.*) infections, use 60 ppm (1200 g DEWORMIT-F 5% per ton of feed) for 7 days.
 - **Turkeys:** 20 ppm Flubendazole (400 g DEWORMIT-F 5% per ton of feed) for 7 days.
 - **Pheasants:** 60 ppm Flubendazole (1200 g DEWORMIT-F 5% per ton of feed) for 7 days.



VETERINARY
NOT FOR HUMAN USE
FOR ANIMAL TREATMENT ONLY



FUREVER 9 CORPORATION
207, Turner Road, Clement Town
Dehradun 248001 (U.K.) India.

ERYCIN

ERYTHROMYCIN THIOCYANATE POWDER FOR ORAL ADMINISTRATION

COMPOSITION

Each gram contains :
Erythromycin thiocyanate 200 mg
Excipients ad 1 g

INDICATIONS

ERCIN is indicated for the treatment of respiratory infections in poultry, including:

- Chronic Respiratory Disease (CRD) caused by *Mycoplasma* spp.
- Infectious diseases caused by erythromycin-sensitive organisms such as:
 - *Staphylococcus aureus* (e.g., infectious arthritis)
 - *Haemophilus paragallinarum* (e.g., infectious coryza)
 - *Pasteurella multocida* (e.g., pasteurellosis, pneumonia)

CONTRAINDICATIONS

Do not administer to:

- Animals with known hypersensitivity to erythromycin
- Animals with severely impaired liver and/or kidney function


DOSAGE AND ADMINISTRATION

For oral administration via drinking water:

- Poultry: 100 g per 100 litres of drinking water for 3–5 consecutive days

WITHDRAWAL PERIOD

- Meat: 3 days
- Eggs: 6 days



VETERINARY
NOT FOR HUMAN USE
FOR ANIMAL TREATMENT ONLY



FUREVER 9 CORPORATION
207, Turner Road, Clement Town
Dehradun 248001 (U.K.) India.

PACK SIZE
Available in
1 Kg Jar
100 g Sachet

EZYDOXIN 200


DOXYCYCLINE HYCLATE

COMPOSITION

Each gram contains :
 Doxycycline hyclate B.P. 200 mg
 Excipients ad 1 g

INDICATIONS

Indicated for the treatment of gastrointestinal and respiratory infections caused by *Doxycycline*-sensitive microorganisms, including: *Bordetella*, *Campylobacter*, *Chlamydia*, *Escherichia coli*, *Haemophilus*, *Mycoplasma*, *Pasteurella*, *Rickettsia*, *Salmonella*, *Staphylococcus*, and *Streptococcus* spp. Applicable for use in **Calves, Goats, Poultry, Sheep, and Swine**.



**VETERINARY
 NOT FOR HUMAN USE
 FOR ANIMAL TREATMENT ONLY**

CONTRAINDICATIONS

- Do not administer to animals with severe hepatic impairment.
- Contraindicated with concurrent use of penicillins, cephalosporins, quinolones, or cycloserine.
- Not to be used in animals with active microbial digestion.
- Hypersensitivity to tetracyclines, penicillins, cephalosporins, quinolones, or cycloserine

DOSAGE AND ADMINISTRATION

For oral administration only:

- **Calves, Goats, and Sheep:** 5 g per 200 kg body weight, twice daily for 3–5 days.
- **Poultry and Swine:** 1 kg per 1000–2000 litres of drinking water for 3–5 days.

Note: For pre-ruminant Calves, Lambs, and Kids only.

WITHDRAWAL PERIOD

- **Meat:**
 - Calves, Goats, Sheep: 14 days
 - Swine: 8 days
 - Poultry: 7 days



FUREVER 9 CORPORATION
 207, Turner Road, Clement Town
 Dehradun 248001 (U.K.) India.

PACK SIZE
 Available in
1 Kg Jar
30 g & 100 g Sachet

EZYDOXIN C

DOXYCYCLINE HYCLATE & COLISTIN SULPHATE

COMPOSITION


Each gram contains:

Doxycycline hyclate	100 mg
Colistin sulphate	1 200 000 IU
Excipients ad	1 g

INDICATIONS

Indicated for the treatment of **gastrointestinal and respiratory infections** caused by microorganisms sensitive to doxycycline and/or colistin, including: *Bordetella*, *Campylobacter*, *Chlamydia*, *Escherichia coli*, *Klebsiella*, *Haemophilus*, *Mycoplasma*, *Pasteurella*, *Rickettsia*, *Salmonella*, *Staphylococcus*, and *Streptococcus spp.*

Recommended for use in **Calves, Goats, Poultry, Sheep, and Swine.**



VETERINARY
NOT FOR HUMAN USE
FOR ANIMAL TREATMENT ONLY

CONTRAINDICATIONS

- Hypersensitivity to tetracyclines and/or polymyxins
- Animals with severe renal or hepatic impairment
- Animals with active microbial digestion (e.g., mature ruminants)
- Animals producing **milk or eggs** intended for human consumption

DOSAGE & ADMINISTRATION

For oral administration:

- **Calves, Goats, Sheep:** 1 g per 20 kg body weight, twice daily for 4–5 days
- **Poultry:** 1 kg per 2000 litres of drinking water for 4–5 days
- **Swine:** 1 kg per 1000 litres of drinking water for 4–5 days

Note: For pre-ruminant calves, lambs, and kids only. Not for use in animals producing milk or eggs for human consumption.

WITHDRAWAL PERIOD

- **Meat:**
 - Calves, Goats, Sheep: 14 days
 - Swine: 8 days ○ Poultry: 7 days



FUREVER 9 CORPORATION
207, Turner Road, Clement Town
Dehradun 248001 (U.K.) India.

PACK SIZE

Available in
1 Kg Jar
100 g Sachet

EZYDOXIN G

GENTAMICIN SULPHATE / DOXYCYCLINE HYCLATE FOR ORAL ADMINISTRATION


COMPOSITION

Each gram contains :

Doxycycline hyclate B.P	100 mg
Gentamicin sulphate B.P	50 mg
Excipients ad	1 g

INDICATIONS

Indicated for the treatment of gastrointestinal and respiratory infections caused by micro-organisms sensitive to doxycycline and/or gentamicin, such as: *Bordetella*, *Campylobacter*, *Chlamydia*, *Escherichia coli*, *Klebsiella*, *Haemophilus*, *Mycoplasma*, *Pasteurella*, *Rickettsia*, *Salmonella*, *Staphylococcus*, and *Streptococcus* spp. Suitable for use in Calves, Goats, Poultry, Sheep, and Swine.



VETERINARY
NOT FOR HUMAN USE
FOR ANIMAL TREATMENT ONLY

CONTRAINDICATIONS

Do not administer in cases of:

- Known hypersensitivity to tetracyclines and/or aminoglycosides
- Severe renal or hepatic impairment
- Concurrent use with penicillins, cephalosporins, quinolones, or cycloserine
- Active microbial digestion

DOSAGE AND ADMINISTRATION

For oral administration:

- **Calves, Goats, and Sheep:** 1 g per 20 kg body weight, twice daily for 3–5 days
- **Poultry and Swine:** 1 kg per 1000 litres of drinking water for 3–5 days

Note: Intended for use in pre-ruminant Calves, Lambs, and Kids only. Not recommended for lactating animals or laying birds.

WITHDRAWAL PERIOD

- **Meat:**
 - Calves, Goats, Sheep: 14 days
 - Swine: 8 days
 - Poultry: 7 days



F9
FUREVER 9 CORPORATION
207, Turner Road, Clement Town
Dehradun 248001 (U.K.) India.



PACK SIZE
Available in
1 Kg Jar
100 g Sachet

GASTROFLU

Flumequine Powder

COMPOSITION

Each gram contains :
 Flumequine 200 mg
 Excipients ad 1 g

INDICATIONS

Flumequine is indicated for the treatment of gastrointestinal, respiratory, and urinary tract infections caused by flumequine-sensitive micro-organisms such as *Campylobacter*, *Escherichia coli*, *Haemophilus*, *Pasteurella*, and *Salmonella* spp. Applicable for use in Calves, Goats, Sheep, Swine, and Poultry.


CONTRAINDICATIONS

- Do not use in animals with known hypersensitivity to flumequine.
- Not suitable for animals with severely impaired hepatic and/or renal function.
- Avoid concurrent use with tetracyclines, chloramphenicol, macrolides, or lincosamides.

DOSAGE & ADMINISTRATION

For oral administration:

- **Calves, Goats, and Sheep:**
 - Administer 10 g per 75–150 kg body weight twice daily for 3–5 days.
 - *Note: Only for use in pre-ruminant calves, lambs, and kids.*
- **Poultry and Swine:**
 - 1 kg per 1000–2000 liters of drinking water for 3–5 days.



**VETERINARY / NOT FOR HUMAN USE
FOR ANIMAL TREATMENT ONLY**

WITHDRAWAL PERIOD

- **Meat:**
 - Calves, Goats, Sheep, and Swine – 8 days
 - Poultry – 3 days



FUREVER 9 CORPORATION
 207, Turner Road, Clement Town
 Dehradun 248001 (U.K.) India.

PACK SIZE
 Available in
 Sachets of 100 g
 Jars of 1000 g

HYDRODEX

DEXAMETHASONE AND HYDROCHLOROTHIAZIDE POWDER FOR ORAL ADMINISTRATION

COMPOSITION

Each gram contains:

Dexamethasone

Sodium Phosphate B.P. 0.25 mg


Hydrochlorothiazide B.P. 75 mg

Excipient 1 g

INDICATION

Indicated for the management of various oedematous conditions, including:

- Generalised congestion and oedema
- Sheath oedema
- Anasarca
- Oedema associated with allergic reactions



VETERINARY
NOT FOR HUMAN USE
FOR ANIMAL TREATMENT ONLY

CONTRAINDICATIONS

Do not use in:

- Pregnant animals
- Animals with active viral infections during the viraemic phase
- Animals diagnosed with diabetes mellitus, congestive heart failure, chronic nephritis, osteoporosis, or glaucoma
- Animals suffering from hepatic encephalopathy
- Cases of severe hypokalaemia
- Animals with known hypersensitivity to any of the active components

DOSAGE AND ADMINISTRATION

For oral administration via drinking water

- **Adult Cattle and Horses:**
 - Day 1: 2 sachets
 - Days 2 and 3: 1 sachet per day(Equivalent to 2 g per 25–50 kg body weight)

WITHDRAWAL PERIOD

- **Meat and offal:** 6 days
- **Milk:** 3 days



F9 CORPORATION
207, Turner Road, Clement Town
Dehradun 248001 (U.K.) India.

PACK SIZE
Available in
20 g Sachet

LEVASAFE

LEVAMISOLE HYDROCHLORIDE

COMPOSITION

Each gram Contains :
Levamisole hydrochloride 200 mg
Excipients ad 1 g

INDICATIONS

For the **prophylaxis and treatment** of gastrointestinal and lungworm infections in:

- **Cattle, Calves, Sheep, and Goats:**


Bunostomum, Chabertia, Cooperia, Dictyocaulus, Haemonchus, Nematodirus, Ostertagia, Protostrongylus, Trichostrongylus spp.

- **Poultry:**

Ascaridia, Capillaria spp.

- **Swine:**

Ascaris suum, Hyostrongylus rubidus, Metastrongylus elongatus, Oesophagostomum spp., *Trichuris suis*



VETERINARY
NOT FOR HUMAN USE
FOR ANIMAL TREATMENT ONLY

CONTRAINDICATIONS

- Do not administer to animals with **impaired hepatic function**.
- Avoid **concurrent administration** with **pyrantel, morantel, or organophosphates**.

DOSAGE AND ADMINISTRATION

For oral administration:

- **Cattle, Calves, Sheep, and Goats:**

7.5 g per 200 kg body weight, single dose

- **Poultry and Swine:**

1 kg per 2000 litres of drinking water, single day treatment

WITHDRAWAL PERIOD

- **Meat:** 10 days
- **Milk:** 4 days



PACK SIZE
Available in
1 Kg Jar
100 g Sachet



FUREVER 9 CORPORATION
207, Turner Road, Clement Town
Dehradun 248001 (U.K.) India.

METROFUR

METRONIDAZOLE 10% Powder

COMPOSITION

Each pack contains :

Metronidazole (as benzoate)	10%
Excipient	q.s.

INDICATIONS

For the treatment of:

- Cankers
 - Giardiasis
 - Other protozoal infections
- Suitable for use in birds.

CONTRAINDICATIONS

- Do not use in animals with **hepatic disorders**.
- Do not use in cases of **hypersensitivity** to metronidazole or any of the excipients.

DOSAGE AND ADMINISTRATION

For oral administration through drinking water:

- **Dosage:** 2 teaspoons per quart of drinking water
- **Duration:** 5 consecutive days
- **Important:** If the bird does not improve or the condition worsens, consult a veterinarian.



FUREVER 9 CORPORATION
207, Turner Road, Clement Town
Dehradun 248001 (U.K.) India.

VETERINARY
NOT FOR HUMAN USE
FOR ANIMAL TREATMENT ONLY



PACK SIZE

Available in **100 g Sachet & 1 Kg Jar**



NEOWIN

Neomycin Sulphate

COMPOSITION

Each gram contains :
Neomycin sulphate 700 mg
Excipients ad 1 g

INDICATIONS

NEOWIN is recommended for the prevention and treatment of bacterial enteritis in calves, sheep, goats, swine, and poultry. It is effective against infections caused by neomycin-sensitive bacteria such as *Escherichia coli*, *Salmonella* spp., and *Campylobacter* spp.


CONTRAINDICATIONS

- Do not use in animals known to be hypersensitive to neomycin.
- Not suitable for animals with severely impaired kidney function.
- Avoid use in animals with active microbial digestion (e.g., ruminants).
- Not to be administered to laying poultry producing eggs intended for human consumption.

DOSAGE & ADMINISTRATION

For oral use only:

- **Calves, Goats, and Sheep:**
 - 10 mg of neomycin sulphate per kg body weight (equivalent to 14 mg/kg of NEOCIN WS) daily for 3–5 days.
 - Note: Only for use in pre-ruminant calves, lambs, and kids.
- **Poultry and Swine:**
 - 300 g per 2000 liters of drinking water for 3–5 days.



**VETERINARY / NOT FOR HUMAN USE
FOR ANIMAL TREATMENT ONLY**

WITHDRAWAL PERIOD

- **Meat:**
 - Calves, Goats, Sheep, and Swine – 21 days
 - Poultry – 7 days



PACK SIZE
Available in
Sachets of 30 g and 100 g
Jars of 1000 g



FUREVER 9 CORPORATION
207, Turner Road, Clement Town
Dehradun 248001 (U.K.) India.

OXYFUR 40%

Oxytetracycline Hydrochloride

COMPOSITION

Each gram contains :
Oxytetracycline hydrochloride B.P. 400 mg
Excipients ad 1 g

INDICATIONS

OXYFUR 40% is indicated for the treatment of gastrointestinal and respiratory tract infections caused by oxytetracycline-sensitive organisms, including *Bordetella*, *Bacillus*, *Corynebacterium*, *Campylobacter*, *E. coli*, *Haemophilus*, *Pasteurella*, *Salmonella*, *Staphylococcus*, *Streptococcus*, *Mycoplasma*, *Rickettsia*, and *Chlamydia* spp.

Recommended for use in Calves, Goats, Sheep, Swine, and Poultry.

CONTRAINDICATIONS

- Do not use in animals with known hypersensitivity to tetracyclines.
- Not recommended in animals with impaired renal or hepatic function.
- Avoid concurrent use with penicillins, cephalosporins, quinolones, or cycloserine.
- Contraindicated in animals with active microbial digestion.


DOSAGE & ADMINISTRATION

For oral administration:

- **Calves, Goats, and Sheep:**
 - 1 g per 20–40 kg body weight, administered twice daily for 3–5 days.
 - *Note: For pre-ruminant calves, lambs, and kids only.*
- **Poultry and Swine:**
 - 1 kg per 2000 liters of drinking water for 3–5 days.



FUREVER 9 CORPORATION
207, Turner Road, Clement Town
Dehradun 248001 (U.K.) India.



**VETERINARY / NOT FOR HUMAN USE
FOR ANIMAL TREATMENT ONLY**

WITHDRAWAL PERIOD

• Meat:

- Calves, Goats, Sheep, and Swine – 8 days
- Poultry – 6 days



PACK SIZE
Available in
Sachets of 30 g and 100 g
Jars of 1000 g

TRIPOWER K3

**Sulphadimidine + Sulphadiazine +
Sulphathiazole + Vitamin K3**

COMPOSITION

Sulphadimidine sodium	B.P.	125 mg
Sulphadiazine sodium	B.P.	125 mg
Sulphathiazole sodium	B.P.	125 mg
Vitamin K3	B.P.	10 mg

INDICATIONS

TRIPOWER K3 is recommended for the treatment of:

- Gastrointestinal diarrhoea
- Respiratory tract infections
- Shipping fever in large animals
- Intestinal coccidiosis in poultry


Warning:

Not to be used in poultry producing eggs intended for human consumption.

DOSAGE & ADMINISTRATION

For oral administration:

- **Poultry:**
 - 1 g of Trisulfacin WS per 2–4 litres of drinking water
 - Dosage schedule: 3 days of treatment, followed by 2 days without treatment, then another 2 days of treatment
- **Calves, Lambs, and Kids:**
 - 1 g per 5 kg of body weight daily for 3–4 days



**VETERINARY / NOT FOR HUMAN USE
FOR ANIMAL TREATMENT ONLY**

WITHDRAWAL PERIOD

- Poultry (meat): 5 days
- Other animals (meat): 21 days



FUREVER 9 CORPORATION
207, Turner Road, Clement Town
Dehradun 248001 (U.K.) India.

PACK SIZE
Available in
Sachets of 30 g and 100 g
Jars of 1000 g

PEZINE

Piperazine Citrate Powder For Oral Administration

COMPOSITION

Each gram contains:
Piperazine citrate B.P. 1000 mg

INDICATIONS

Piperazine Citrate is effective for the treatment of nematode infections in horses, swine, cattle, and poultry.

- **Swine:** Effective against *Oesophagostomum* and *Ascaris*.
- **Cattle:** Effective against *Ascaris*, *Nematodirus*, *Ostertagia*, *Cooperia*, and *Oesophagostomum*.
- **Horses:** Effective against *Strongylus*, *Oxyuris*, *Trichonema*, and *Ascaris*.
- **Poultry:** Effective against *Ascaris* and *Capillaria*.


CONTRAINDICATIONS

- Do not administer to animals with known hypersensitivity to piperazine.
- Not recommended for animals with severely impaired liver and/or kidney function.

DOSAGE & ADMINISTRATION

For oral use only:

- **Horses, Swine, and Cattle:** 1–2 g per 10 kg body weight, administered via drinking water.
- **Poultry:** 1 kg per 1000 liters of drinking water for 2 consecutive days.



**VETERINARY / NOT FOR HUMAN USE
FOR ANIMAL TREATMENT ONLY**

WITHDRAWAL PERIOD

- **Meat:** 2 days
- **Eggs:** 2 days



FUREVER 9 CORPORATION
207, Turner Road, Clement Town
Dehradun 248001 (U.K.) India.

PACK SIZE
Available in
Sachets of 100 g
Jars of 1000 g

TYLODOX

Doxycycline Hyclate / Tylosin Tartrate
Powder For Oral Administration

COMPOSITION

Each gram contains :
Doxycycline hyclate B.P. 100 mg
Tylosin tartrate B.P. 100 mg
Excipients ad 1 g

INDICATIONS

TYLODOX is indicated for the treatment of gastrointestinal and respiratory tract infections caused by microorganisms sensitive to doxycycline and tylosin. Target pathogens include *Bordetella*, *Campylobacter*, *Chlamydia*, *E. coli*, *Haemophilus*, *Mycoplasma*, *Pasteurella*, *Rickettsia*, *Salmonella*, *Staphylococcus*, *Streptococcus*, and *Treponema* species. Recommended for use in Calves, Goats, Sheep, Swine, and Poultry.


CONTRAINDICATIONS

- Do not administer to animals with severe hepatic impairment.
- Avoid concurrent use with penicillins, cephalosporins, quinolones, or cycloserine.
- Not recommended for animals known to be hypersensitive to tetracyclines and/or tylosin.
- Contraindicated in animals with active microbial digestion.

DOSAGE & ADMINISTRATION

For oral administration:

- **Calves, Goats, and Sheep:**
 - 5 g per 100 kg body weight, administered twice daily for 3 to 5 days.
 - *Note: For pre-ruminant calves, lambs, and kids only.*
- **Poultry and Swine:**
 - 1 kg per 1000–2000 liters of drinking water for 3 to 5 days.



**VETERINARY / NOT FOR HUMAN USE
FOR ANIMAL TREATMENT ONLY**

WITHDRAWAL PERIOD

- **Meat:**
 - Calves, Goats, and Sheep – 14 days
 - Swine – 8 days
 - Poultry – 7 days



FUREVER 9 CORPORATION
207, Turner Road, Clement Town
Dehradun 248001 (U.K.) India.

PACK SIZE
Available in
Sachets of 100 g
Jars of 1000 g

TYLOSOL

TYLOSIN TARTRATE

COMPOSITION


Each gram contains :
Tylosin tartrate B.P. 1000 mg

INDICATIONS

Indicated for the treatment of gastrointestinal and respiratory infections caused by *Tylosin*-sensitive microorganisms, including:

Campylobacter, *Mycoplasma*,
Pasteurella, *Staphylococcus*, *Streptococcus*, and
Treponema spp.

Applicable for use in **Calves, Goats, Poultry, Sheep,**
and **Swine**.



VETERINARY
NOT FOR HUMAN USE
FOR ANIMAL TREATMENT ONLY

CONTRAINDICATIONS

- Hypersensitivity to tylosin.
- Do not administer concurrently with penicillins, cephalosporins, quinolones, or cycloserine.
- Not for animals with active microbial digestion.

DOSAGE AND ADMINISTRATION

For oral administration only:

- **Calves, Goats, and Sheep:** 5 g per 220–250 kg body weight, twice daily for 5–7 days.
- **Poultry:** 1 kg per 1500–2000 litres of drinking water for 3–5 days.
- **Swine:** 1 kg per 3000–4000 litres of drinking water for 5–7 days.

WITHDRAWAL PERIOD

- **Meat:**
 - Calves, Goats, Poultry, Sheep: 5 days
 - Swine: 3 days



PACK SIZE
Available in
1 Kg Jar
30 g & 100 g Sachet



FUREVER 9 CORPORATION
207, Turner Road, Clement Town
Dehradun 248001 (U.K.) India.